

Petlife

cimicat™



THE NATURAL FOOD FOR CATS AND KITTENS

Cimicat is scientifically prepared food for cats of all ages. It helps to give a smooth sleek coat and promotes good health.

- ♥ Nourishes cats of all ages.
- ♥ Ensures proper rearing of orphan kittens or those in large litter, and later as a food supplement.
- ♥ Provides the pregnant and lactating queen with the essential substances for nutrition and growth.



Petlife

Petlife International Limited, Unit 2, Cavendish Road, Bury St Edmunds, Suffolk IP33 3TE
t. 01284 761131 f. 01284 761139 e. info@petlifeonline.co.uk w. www.petlifeonline.co.uk

Mixing Instructions

Boil water and allow to cool to maximum 40°C (lukewarm/tepid). Mix at a rate of 20% dry matter i.e. 1 part powder to 2 parts water: Add a small amount of water and mix to an even paste, then add the remaining required water, stirring all the time.

1 x 10ml scoop (5gm) Cimicat powder to 20ml (20gm) water.

To make 25ml: Add 1 x 10ml scoop of Cimicat powder to 2 x 10ml scoops of water.

DAILY FEEDING GUIDE

Age	Prepared liquid Cimicat at recommended mixing rate	Number of feeds per day	Liquid Cimicat per feed
1 – 3 days	20ml - 25ml	Every 2 hours	2ml - 2.5ml
4 - 7 days	30ml - 35ml	10 to 12 times daily	3ml - 3.5ml
8 - 10 days	35ml - 40ml	Every 2 hours / 10 times daily	3.5ml - 4ml
11 - 14 days	40ml - 50ml	Every 3 hours	5ml - 6.25ml
15 - 21 days	50ml - 75ml	Every 4 hours	8.25ml - 12.5ml
8 - 14 days	40ml - 100ml	Every 3 hours / 8 times daily	5ml - 12.5ml
15 - 21 days	75ml - 135ml	Every 3 - 4 hours / 6 times daily	12.5ml - 22.5ml
21 days up to 6 weeks	50ml - 100ml in addition to normal feed	3 - 4 times daily	12.5ml - 25ml

The above is merely a guide and feeding rates will depend on the weight of the kitten and other circumstances. As a general guide feed 20ml - 25ml of mixed Cimicat liquid per 100gm of kitten weight. Caution: Both overfeeding and underfeeding can cause serious health problems with your kitten. Kittens have small immature kidneys, which cannot cope with excessive amounts of liquid.

Overfeeding can cause diarrhoea, whilst underfeeding can cause dehydration.

